JOURNAL OF CHROMATOGRAPHY B

VOL. 698 NOS. 1 + 2 26 SEPTEMBER 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

Regular Papers

Preparation and gas chromatographic–mass spectrometric analysis of N-acetyl-O-trimethylsilyl derivatives of long-chain base residues of natural sphingomyelin
by N.U. Olsson, P. Kaufmann and S. Dzeletović (Stockholm, Sweden) (Received 15 May 1997) 1

Reversed-phase high-performance liquid chromatographic method for the determination of the 11-hydroxythromboxane B_2 anomers equilibrium
by H. John and W. Schlegel (Münster, Germany) (Received 12 May 1997) 9

Application of [^{125}I]-[Tyr^8]-substance P prepared by the chloramine-T method to receptor-binding experiments after subsequent reduction with mercaptoethanol and purification by reversed-phase liquid chromatography
by K. Rissler and H. Cramer (Freiburg, Germany) and P. Engelmann (Herbolzheim, Germany) (Received 18 April 1997) . . . 17

Alteration of redox state of human serum albumin in patients under anesthesia and invasive surgery
by A. Hayakawa, K. Kuwata, S. Era, M. Sogami, H. Shimonaka, M. Yamamoto, S. Dohi and H. Hirose (Gifu, Japan) (Received 29 April 1997) 27

Use of immobilized metal ion affinity chromatography and dye–ligand chromatography for the separation and purification of rainbow trout pituitary gonadotropins, GTH I and GTH II
by M.S. Govoroun (Rennes, France), J.-C. Huet and J.-C. Pernollet (Jouy en Josas, France) and B. Breton (Rennes, France) (Received 18 April 1997) 35

Effect of different surfactants on the separation by micellar electrokinetic chromatography of a complex mixture of dipeptides in urine of prolidase-deficient patients
by R. Grimm (Waldbronn, Germany) and G. Zanaboni, S. Viglio, K.M. Dyne, G. Cetta and P. Iadarola (Pavia, Italy) (Received 21 May 1997) 47

Fractionation of RNA polymerase II transcription factors from Hela cell nuclear extracts by affinity chromatography on "DNA-like" phosphorylated polystyrene
by E. Imbert, D. Letourneur and M. Jozefowicz (Villetaneuse, France) (Received 7 April 1997) 59

High-performance liquid chromatographic separation and tandem mass spectrometric identification of breakdown products associated with the biological hydrolysis of a biomedical polyurethane
by G.B. Wang and J.P. Santerre (Toronto, Canada) and R.S. Labow (Ottawa, Canada) (Received 1 May 1997) 69

Determination of the main hydrolysis products of organophosphorus nerve agents, methylphosphonic acids, in human serum by indirect photometric detection ion chromatography
by M. Katagi, M. Nishikawa, M. Tatsuno and H. Tsuchihashi (Osaka, Japan) (Received 18 April 1997) 81

Dye-incorporated poly(EGDMA–HEMA) microspheres as specific sorbents for aluminum removal
by A. Denizli, B. Salih, C. Kavakli and E. Pişkin (Ankara, Turkey) (Received 12 May 1997) 89

Simple and rapid determination of enflurane in human tissues using gas chromatography and gas chromatography–mass spectrometry
by H. Ise, K. Kudo, N. Jitsufuchi, T. Imamura and N. Ikeda (Fukuoka, Japan) (Received 14 April 1997) 97

(Continued overleaf)

Contents (continued)

Determination of *p*-trifluoromethylphenol, a metabolite of fluoxetine, in tissues and body fluids using an electron-capture gas chromatographic procedure
by L.J. Urichuk, L.J. Aspeslet, A. Holt, P.H. Silverstone, R.T. Coutts and G.B. Baker (Edmonton, Canada) (Received 1 May 1997) 103

Determination of *N,N′,N″*-triethylenthiophosphoramide in biological samples using capillary gas chromatography
by R.J. van Maanen, R.D. van Ooijen and J.H. Beijnen (Utrecht, The Netherlands) (Received 14 April 1997) 111

Quantitative determination of BMS-186716, a thiol compound, in rat plasma by high-performance liquid chromatography–positive ion electrospray mass spectrometry after hydrolysis of the methyl acrylate adduct by the native esterases
by M. Jemal and D.J. Hawthorne (New Brunswick, NJ, USA) (Received 22 April 1997) 123

Buspirone metabolite structure profile using a standard liquid chromatographic–mass spectrometric protocol
by E.H. Kerns (New Brunswick, NJ, USA), R.A. Rourick and K.J. Volk (Wallingford, CT, USA) and M.S. Lee (Princeton, NJ, USA) (Received 14 April 1997) 133

Highly sensitive high-performance liquid chromatographic–tandem mass spectrometric method for the analysis of dextromethorphan in human plasma
by T.H. Eichhold, M. Quijano, W.L. Seibel, C.A. Cruze, R.L.M. Dobson and K. R. Wehmeyer (Mason, OH, USA) (Received 21 May 1997) 147

Determination of cloxacillin in milk and blood of dairy cows by high-performance liquid chromatography
by B. Pérez and C. Prats (Bellaterra, Spain), E. Castells (Barcelona, Spain) and M. Arboix (Bellaterra, Spain) (Received 28 April 1997) 155

Automated determination of tramadol enantiomers in human plasma using solid-phase extraction in combination with chiral liquid chromatography
by A. Ceccato, P. Chiap, Ph. Hubert and J. Crommen (Liège, Belgium) (Received 23 April 1997) 161

High-performance liquid chromatographic assay for CI-1004, a dual-inhibitor antiinflammatory agent, in rat, rabbit, dog, monkey and human plasma
by L.D. Andress and R.J. Guttendorf (Ann Arbor, MI, USA) (Received 18 April 1997) 171

Enantioselective high-performance liquid chromatographic determination of nicardipine in human plasma
by T. Uno, T. Ohkubo and K. Sugawara (Hirosaki, Japan) (Received 21 April 1997) 181

Simultaneous determination of atenolol and chlorthalidone in plasma by high-performance liquid chromatography. Application to pharmacokinetic studies in man
by C. Giachetti, A. Tenconi, S. Canali and G. Zanolo (Ivrea, Italy) (Received 22 April 1997) 187

Isolation of N,N-dialkylated derivatives of valproylglycinamide from dog plasma by active charcoal adsorption and their quantification by high-performance liquid chromatography
by O. Spiegelstein, M. Bialer and B. Yagen (Jerusalem, Israel) (Received 10 April 1997) 195

Automated determination of levodopa and carbidopa in plasma by high-performance liquid chromatography–electrochemical detection using an on-line flow injection analysis sample pretreatment unit
by A. Tolokán, I. Klebovich, K. Balogh-Nemes and G. Horvai (Budapest, Hungary) (Received 28 April 1997) 201

Simplified high-performance liquid chromatographic method for determination of risperidone and 9-hydroxyrisperidone in serum from patients comedicated with other psychotropic drugs
by O.V. Olesen and K. Linnet (Risskov, Denmark) (Received 12 May 1997) 209

Simultaneous determination of pyrimethamine and sulphadoxine in human plasma by high-performance liquid chromatography after automated liquid–solid extraction
by H. Astier, C. Renard, V. Cheminel, O. Soares, C. Mounier, F. Peyron and J.F. Chaulet (Lyon, France) (Received 16 April 1997) 217

Prevalidation statistical design to assess analytical methods. Example of a quick liquid chromatographic assay of itraconazole in serum
by M. Cociglio, D. Hillaire-Buys and R. Alric (Montpellier, France) (Received 18 April 1997) 225

Determination of saquinavir in human plasma, saliva, and cerebrospinal fluid by ion-pair high-performance liquid chromatography with ultraviolet detection
by R.M.W. Hoetelmans, M. van Essenberg, P.L. Meenhorst, J.W. Mulder and J.H. Beijnen (Amsterdam, The Netherlands) (Received 4 April 1997) 235

Contents (continued)

Rapid and simple method to determine chloroquine and its desethylated metabolites in human microsomes by high-performance liquid chromatography with fluorescence detection
by J. Ducharme and R. Farinotti (Chatenay-Malabry, France) (Received 10 April 1997) 243

High-performance liquid chromatographic determination of oxolinic acid in the plasma of seabass (*Dicentrarchus labrax*) anaesthetized with 2-phenoxyethanol
by S. Loussouarn, H. Pouliquen and F. Armand (Nantes, France) (Received 19 April 1997) 251

Sensitive and convenient high-performance liquid chromatographic method for the determination of mitomycin C in human plasma
by G. Joseph, W. Biederbick, U. Woschée, M. Theisohn and W. Klaus (Cologne, Germany) (Received 23 April 1997) 261

Solid-phase extraction and high-performance liquid chromatographic determination of tamoxifen and its major metabolites in breast tumour tissues
by J. MacCallum, J. Cummings, J.M. Dixon and W.R. Miller (Edinburgh, UK) (Received 24 March 1997) 269

Determination of irinotecan (CPT-11) and its active metabolite SN-38 in human plasma by reversed-phase high-performance liquid chromatography with fluorescence detection
by P. de Bruijn, J. Verweij, W.J. Loos, K. Nooter, G. Stoter and A. Sparreboom (Rotterdam, The Netherlands) (Received 25 April 1997) 277

Determination of idarubicin and idarubicinol in plasma by capillary electrophoresis
by G. Hempel, S. Haberland, P. Schulze-Westhoff, N. Möhling, G. Blaschke and J. Boos (Münster, Germany) (Received 2 May 1997) 287

Simultaneous determination of mycophenolic acid and its glucuronide conjugate in human plasma by a single-run ion-paring method
by J.J.Z. Huang (Palo Alto, CA, USA), H. Kiang (Sunnyvale, CA, USA) and T.L. Tarnowski (Palo Alto, CA, USA) (Received 17 April 1997) 293

Short Communications

Simultaneous determination of urinary cystathionine, lanthionine, *S*-(2-aminoethyl)-L-cysteine and their cyclic compounds using liquid chromatography–mass spectrometry with atmospheric pressure chemical ionization
by S. Yu, K. Sugahara, J. Zhang, T. Ageta and H. Kodama (Kochi, Japan), M. Fontana (Rome, Italy) and S. Duprè (Roma, Italy) (Received 12 May 1997) 301

Determination of purine nucleoside phosphorylase activity in human erythrocytes by capillary electrophoresis
by T. Adam and J. Ševčík (Olomouc, Czech Republic), L.D. Fairbanks (London, UK) and P. Barták (Olomouc, Czech Republic) (Received 12 May 1997) 308

Contribution to the determination of the chemotherapeutic drug G1 in biological fluids
by L. Gil del Valle, D. Gebru W/Michael, A. Martinez Rodríguez, R. Tápanes Peraza, F. Ramos and J. Pérez Avila (Ciudad de la Habana, Cuba) (Received 27 March 1997) 312

Sensitive high-performance liquid chromatographic assay method for the determination of picroside I in plasma
by A.K. Dwivedi, D. Kulkarni and S. Singh (Lucknow, India) (Received 9 April 1997) 317

Related Articles J. Chromatography A. Vol. 774 321

Author Index Volume 698 322

Compound Index Volume 698 325

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc